

## MOTIVATION

- ❖ Scalability in Bayesian nonparametric.
- ❖ Hard clustering for fast inference.
- ❖ Small variance asymptotic
- ❖ Multinomial observations with large number of trials.
- ❖ Let assume the number of trials go to infinity.  $\infty$

## CONTRIBUTIONS

- ❖ A novel view of Large Sample Asymptotic (LSA).
- ❖ Theoretical derivation for the proposed LSA on Dirichlet Process Mixture model with count data.
- ❖ Comparing with the existing clustering techniques.

## Dirichlet Process Mixture - Gibbs Sampler

Gibbs sampler for posterior inference

$$p(z_i = k | z_{-i}, x_i, \phi, \alpha, G) \propto \begin{cases} N_k \times p(x_i | \phi_{z_i}) & \text{used } k \\ \alpha \times \int_{\phi} p(x_i | \phi) dG(\phi) & \text{new } k \end{cases}$$



## DPM - Small Variance Asymptotic

Assume the data variance approaches to zero  $\sigma \rightarrow \infty$

$$\lim_{\sigma \rightarrow 0} \hat{\gamma}(z_i = k) = \begin{cases} D_{\phi}(x_i, \mu_k) & \text{used } k \\ \lambda & \text{new } k \end{cases}$$

Where  $D_{\phi}(x_i, \mu_k)$  is a Bregman divergence.



## Large Sample Asymptotic

Multinomial likelihood  $p(x | \phi) = \frac{n!}{\prod_{d=1}^D x_d!} \prod_{d=1}^D \phi_d^{x_d}$

Average Log Likelihood  $\bar{L} \equiv \log p(x | \phi)^{\frac{1}{n}}$

Equivalently  $\bar{L} = \frac{1}{n} \log n! - \frac{1}{n} \sum_{d=1}^D \log x_d! + \sum_{d=1}^D \frac{x_d}{n} \log \phi_d$

Stirling Approximation  $\log n! = n \log n - n + O(\log n)$

Alternative representation of Multinomial likelihood

$$P(x | \phi) = \exp\{-n D_{KL}(\hat{x} | \phi) + O(\log n) + \sum_{d=1}^D O(\log x_d)\}$$

## DPM - LSA

Sampling  $z_i$

$$p(z_i = k) \propto N_k \times \exp[-n D_{KL}(\hat{x}_i | \phi_{z_i}) + T]$$

Sampling  $z_i^{\text{new}}$

$$p(z_i = k^{\text{new}}) \propto p(z_i = k^{\text{new}} | z_{-i}, \alpha) p(x_i | z_i = k^{\text{new}}, G)$$

$$p(z_i = k^{\text{new}}) \propto \alpha \frac{n!}{\prod_{d=1}^D x_{id}!} \frac{\Gamma(\sum_{d=1}^D \gamma_d)}{\Gamma(\sum_{d=1}^D [\gamma_d + x_{id}])} \prod_{d=1}^D \frac{\Gamma(x_{id} + \gamma_d)}{\Gamma(\gamma_d)}$$

$$p(z_i = k^{\text{new}}) \propto C(x_i) \exp(-n\lambda) \exp[O(\log n)]$$

Let the number of trials in Multinomial distribution go to infinity

$$\lim_{n \rightarrow \infty} \hat{\gamma}(z_i = k) = \begin{cases} D_{KL}(\hat{x}_i | \phi_k) & \text{used } k \\ \lambda & \text{new } k. \end{cases}$$



## Experiments

	Approach	AP	K-means	GMM	DPM	DPmeans	DPM-LSA
NUS WIDE dataset • 3411 images • SIFT features (500 dimensions) Matlab environment	Setting		Euclidian distance		Collapsed Gibbs	$\lambda = 5960$	$\lambda = 1.73$
	#Cluster	K=18	K=5-30	K=5-30	K=17	K=17	K=19
	NMI	0.166	0.19(.01)	0.19(.01)	<b>0.188</b>	0.161	0.174
	F1score	0.145	0.16(.01)	0.16(.01)	0.173	0.166	<b>0.184</b>
	Time	167.8	14	15	1200	38	39.8